

## **Associations between resilience and the well-being of mothers of children with autism spectrum disorder and other developmental disabilities**

Halstead, Elizabeth; Ekas, Naomi; Hastings, Richard; Griffith, Gemma

### **Journal of Autism and Developmental Disorders**

DOI:

[10.1007/s10803-017-3447-z](https://doi.org/10.1007/s10803-017-3447-z)

Published: 01/04/2018

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Halstead, E., Ekas, N., Hastings, R., & Griffith, G. (2018). Associations between resilience and the well-being of mothers of children with autism spectrum disorder and other developmental disabilities. *Journal of Autism and Developmental Disorders*, 48(4), 1108-1121.  
<https://doi.org/10.1007/s10803-017-3447-z>

#### **Hawliau Cyffredinol / General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Associations between Resilience and the Well-Being of Mothers of Children with Autism  
Spectrum Disorder and Other Developmental Disabilities

Elizabeth Halstead  
Bangor University

Naomi Ekas  
Texas Christian University

Richard P. Hastings  
University of Warwick

Gemma M. Griffith  
Bangor University

Corresponding author information: Naomi Ekas, Texas Christian University,  
[naomi.ekas@tcu.edu](mailto:naomi.ekas@tcu.edu), 817-257-7848

### Abstract

There is variability in the extent to which mothers are affected by the behavior problems of their children with developmental disabilities (DD). We explore whether maternal resilience functions as a protective or compensatory factor. In Studies 1 and 2, using moderated multiple regression models, we found evidence that maternal resilience functioned as a compensatory factor – having a significant independent main effect relationship with well-being outcomes in mothers of children with DD and autism spectrum disorder. However, there was no longitudinal association between resilience and maternal well-being outcomes. There was little evidence of the role of resilience as a protective factor between child behavior problems and maternal well-being in both studies.

*Key words:* autism spectrum disorder, developmental disability, mothers, psychological well-being, resilience

## Abstract

There is variability in the extent to which mothers are affected by the behavior problems of their children with developmental disabilities (DD). We explore whether maternal resilience functions as a protective or compensatory factor. In Studies 1 and 2, using moderated multiple regression models, we found evidence that maternal resilience functioned as a compensatory factor – having a significant independent main effect relationship with well-being outcomes in mothers of children with DD and autism spectrum disorder. However, there was no longitudinal association between resilience and maternal well-being outcomes. There was little evidence of the role of resilience as a protective factor between child behavior problems and maternal well-being in both studies.

*Key words:* autism spectrum disorder, developmental disability, mothers, psychological well-being, resilience

## Associations between Resilience and the Well-Being of Mothers of Children with Autism

### Spectrum Disorder and Other Developmental Disabilities

Research showing higher levels of stress and other negative psychological well-being outcomes (e.g., depression, anxiety) in mothers of children with developmental disabilities (including autism spectrum disorder and/or intellectual disability) compared to mothers of typically developing children is well established (Hastings, 2016). In addition, research has identified that parents of children with autism spectrum disorder (ASD) experience higher stress levels than parents of children diagnosed with Down's Syndrome, Cerebral Palsy, and Global Developmental Delay (e.g., Blacher & McIntyre, 2006; Dabrowska & Pisula, 2010; Hayes, & Watson, 2013). However, there is variability in how mothers respond to the stress of raising a child with developmental disabilities (DD). In analysis of a population-based sample, for example, Totsika, Hastings, Emerson, Lancaster and Berridge (2011) found that 40% of mothers of children with DD reported experiencing concerning clinical levels of emotional problems, and 60% did not.

Within the context of ASD, the behavioral symptoms of ASD and the severity of symptoms have been explored in association with parental outcomes. It was noted by Smith et al. (2008) that the literature explores mostly associations between child ASD symptoms, specifically, and parental outcomes (e.g., Eisenhower, Baker, & Blancher, 2005; Ello & Donovan, 2005; Lecavalier, Leone, & Wiltz, 2006). However, the question remains if parental stress is primarily due to the child's ASD symptoms or child behavioral and emotional problems more generally (e.g., Hastings, Kovshoff, Ward et al. 2005; Herring et al., 2006).

Previous research has shown behavior problems exhibited by children with DD explain some of the variation in maternal outcomes. Indeed, child behavior problems are a risk factor for lower levels of maternal psychological well-being; having been identified in

several longitudinal studies as a significant predictor of an increase in maternal negative psychological well-being over time (e.g., Baker et al., 2003; Herring et al., 2006; Lecavalier, Leone, & Wiltz, 2006; Zeedyk & Blacher, 2015). Firth and Dryer (2013) also found that children with ASD's behavioral and emotional problems affected overall levels of parental distress, such as stress, tension, anxiety, and depression. The severity of the child's ASD symptoms are also associated with parental stress and depression, suggesting that the severity of the child's ASD symptoms is positively related to the level of parental stress (e.g., Eisenhower, Baker, & Blacher, 2005; Hastings & Johnson, 2001; Hastings et al., 2005; Hill-Chapman et al., 2013). Other aspects of ASD symptomology have also been explored, such as the severity of social impairment, which was found to predict parenting stress (Firth & Dryer, 2013).

Despite the relationship between child behavioral and emotional problems and maternal well-being consistently shown in existing research, there is still variability in mothers' well-being. Not all mothers whose child has significant behavior problems or elevated ASD symptom severity report increased psychological distress or lower levels of well-being. For example, positive parental outcomes are also associated with raising a child with ASD, such as personal growth, improved relationships with others, greater patience, and more empathy (Hastings & Taunt, 2002; Pakenham, Sofronoff, & Samios, 2005; Scorgie & Sobsey, 2000). This variability in mothers' well-being suggests that there are additional factors affecting the relationship between child behavioral problems and maternal well-being. For example, MacDonald, Hastings, and Fitzsimons (2010) found psychological acceptance partially mediated the impact of child behavior problems on paternal stress, anxiety, and depression in a cross-sectional study. In addition, Weiss, Cappadocia, MacMullin, Viecili, and Lunsky (2012) found supporting evidence, also in a cross-sectional study, that maternal empowerment is a partial mediator between child behavior problems and greater maternal

1 distress in mothers of children with ASD. Empowerment is defined as a psychological  
2 process in which an individual is active in changing or eliminating potentially stressful events  
3 through applying knowledge and skill (Gutiérrez, 1994). Self-efficacy was also found to  
4 moderate the effect of child behavior problems on anxiety in fathers of children with ASD  
5 (Hastings & Brown, 2002).  
6  
7  
8  
9  
10

11 One construct which could be important as an explanatory variable for the variability  
12 of maternal well-being in families of children with DD, including ASD, is resilience.  
13 Resilience is of growing interest in mainstream research. For example, Fletcher and Sarkar  
14 (2013) recently reviewed the resilience literature and critiqued the variety of definitions,  
15 concepts and theories of psychological resilience. In the disability field, McConnell and  
16 Savage (2015) proposed expanding the current research agenda to consider the resilience, and  
17 thus adaptation, of families caring for children with DD through greater understanding of the  
18 resources needed to meet everyday challenges. In terms of research addressing resilience in  
19 parents of children with DD, several potential resilience factors have been explored. These  
20 include hope (Lloyd & Hastings, 2009) and self-efficacy (Hastings & Brown, 2002). More  
21 recently, a systematic review by Peer and Hillman (2014) suggested that coping style,  
22 optimism, and social support may all be factors that influence resilience in parents of children  
23 with DD.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 Although some research exists on the resilience of parents of children with DD, there  
44 is a lack of conceptual clarity regarding its definition (Peer & Hillman, 2014). Rutter (1987)  
45 suggested that “resilience is concerned with individual variations in response to risk. Some  
46 people succumb to stress and adversity whereas others overcome life hazards” (p. 317).  
47 Resilience is also defined as “the ability to withstand hardship and rebound from adversity,  
48 becoming more strengthened and resourceful” (Walsh, 1998, 2006, p. 263). These two  
49 definitions are based around a risk/stress – resilience framework: for resilience to be  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 displayed, a stressor must be experienced. Therefore, in the current context resilience might  
2 be demonstrated when mothers report good levels of well-being despite raising a child with  
3 DD who has high levels of behavior problems.  
4  
5

6  
7 Ruiz-Robledillo et al. (2014) found resilience showed associations with overall  
8 general physical and psychological health of caregivers of children with ASD. Research  
9 suggests that resilience is associated with mental health, such as anxiety, insomnia and  
10 depression in caregivers (e.g., Tang et al., 2013). The Resilience Scale for Adults (RSA:  
11 Friborg et al., 2006) has been used in a small population of parents of children with DD, as  
12 part of a comparison between parents of children with Intellectual Disabilities and parents of  
13 children with Sanfilippo syndrome and it was found parents rated *social resources* as their  
14 highest protective factor, and *planned future* as their lowest protective factor (Grant et al.,  
15 2013). Resilience research to date in the DD field is generally based on cross-sectional  
16 research. However, Bayat (2007) suggested that resilience is a process that can only be shown  
17 over time and suggested the need for longitudinal studies. We found only one longitudinal  
18 study with parents of children with DD. This study was by Gertstein, Crnic, Blacher and  
19 Baker (2009), who conducted a longitudinal study exploring the trajectories of daily  
20 parenting stress in parents of young children with DD. The study found factors such as  
21 psychological well-being, marital adjustment and positive parent-child relationships affected  
22 parenting stress differently in mothers and fathers, and it was concluded that parents affect  
23 each other's resilience.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 In the broader literature on resilience, there are three main theoretical ways to  
49 consider resilience: as a compensatory factor (resilience factors have a direct main effect,  
50 reducing negative outcomes directly), as a protective factor (reducing negative outcomes in  
51 the context of exposure to risk – a moderated effect), and finally the challenge model, which  
52 suggests that when exposed to low levels of risk, resilience builds over time (Brook,  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



Whiteman, Gordon, & Cohen, 1986; Brook, Whiteman, Gordon, & Cohen 1989; Fergus & Zimmerman, 2005). In terms of the challenge model, Andrews, Page, and Neilson (1993) suggest that childhood adversities may protect against the effects of later life stress, as this produces “steeling effects” (Lyons & Parker, 2007; Oldehinkel & Ormel, 2014; Rutter, 2006; Seery, Holman, & Silver, 2010).

Each of these different conceptualisations of resilience leads to different predictions about maternal outcomes in DD research. To address the limitations in existing research and to ground the examination of maternal resilience in alternative theoretical models, we conducted two separate studies to examine resilience in mothers of children with DD, including ASD, in two different countries: the UK and the USA. In Study 1 we conceptualised child behavioral and emotional problems as a risk factor likely to lead to lower maternal well-being (stress, mental health status, positive perceptions of raising a child with ASD and other DDs, and perceptions of family satisfaction). If maternal resilience acted as a protective factor, we would expect maternal well-being to be less affected when exposed to high levels of child behavioral and emotional problems if they also score high on a measure of resilience, meaning resilience is affecting maternal outcomes at high levels of risk (high levels of child behavioral problems). If maternal resilience acted as a compensatory factor, we would expect resilience to emerge as a significant independent predictor of maternal outcomes – an effect that is not interactive with child behavioral and emotional problems as a risk factor.

Study 2 was the first to explore resilience in mothers of children with ASD longitudinally, as well as with cross-sectional data. First, this study sought to examine the same cross-sectional associations from Study 1 in a new USA sample of mothers of children with ASD, and with a different maternal resilience measure. This study aimed to determine whether the severity of the child’s ASD symptoms is associated with positive and negative

maternal well-being outcomes when resilience is a moderator, and child ASD symptoms are accounted for. Second, we examined if resilience functions as a compensatory factor affecting maternal well-being outcomes. Third, we explored further how resilience functions longitudinally, thus investigating whether maternal resilience predicts maternal well-being over time.

### Study 1

This first study examined whether maternal resilience served as a compensatory (direct, positive impact on maternal well-being) or protective factor (moderated the association between child behavioral and emotional problems and maternal well-being) among mothers of children with DD residing in the UK.

### Method

**Participants.** The participants were 312 mothers (300 biological mothers, nine adoptive mothers, and three foster mothers) of children aged between four and 15 years old ( $M = 10.02$ ,  $SD = 3.08$ ) with DD, 308 of whom reported they were the primary carer of their child. The mothers' ages ranged from 23 to 67 years ( $M = 42.50$ ,  $SD = 7.13$ ) and 252 were currently living with a spouse or partner. Maternal and child demographic information are shown in Tables 1 and 2, respectively.

**Procedure.** We received approval from an institutional research ethics review board and an external National Independent Research Ethics Committee and local Research and Development offices that are part of the National Health Service (NHS) in the UK. Participants were recruited to complete an online survey through a multi-point recruitment method, which included emailing online links, distributing flyers and information sheets to General Practice (GP) surgeries and secondary care services whose focus was to provide a service for children with DD, UK charities relevant to children with DD, and DD parent support groups. Special Educational Needs schools in North Wales and the North West of

England were sent flyers and information sheets to distribute to parents. Online recruitment via social media (Twitter and Facebook) and online blogs was also on-going throughout the recruitment period. Several participants requested hard copies of the survey and returned completed surveys by post. As all mothers completed all questions, there were no missing data in this dataset. In total, 355 parents responded to the survey. Of the 326 mothers who completed the survey, nine were excluded as their child was not aged between four and 16, and three were excluded because their children did not live with them full time. The fathers who participated in the survey were excluded from this analysis due to the differences seen in previous DD research between mothers and fathers (e.g., Jones, Totsika, Hastings, & Petalas, 2013). Furthermore, it was unknown if the fathers came from the same family as the mothers. Due to the nature of the recruitment methods, we are unable to determine the overall response rate for this survey.

**Measures.** Six measures plus a demographic questionnaire were used in this study; all measures were completed by the mother of the child with IDD.

*Demographic Questionnaire.* Demographic information was gathered using a questionnaire developed by the research team and included questions about the mother (see Table 1 for details) and their child with DD (see Table 2). Demographic variables were re-coded dichotomously: children's physical and sensory abilities, which originally had three categories, were reduced to two, (e.g., able to walk, able to walk with help, or unable to walk without help was reduced to: able to walk with or without help, or unable to walk), maternal employment was re-coded into two categories (no paid employment vs. employed), ethnicity was coded as white British versus all other categories, and education was coded into degree level and above versus lower than degree level. All other demographic variables were dichotomously coded (male vs. female; with child diagnosis in four groups that were dummy coded: autism present vs. no autism present, Down's Syndrome present vs. no Down's

Syndrome present, Cerebral Palsy present vs. no Cerebral Palsy present, and the remainder of children as a mixed IDD group). Socio-Economic Position (SEP) has been associated with maternal well-being in several studies (e.g., Totsika et al., 2011), and so we gathered relevant data and combined several indicators into an index of deprivation. The first indicator was neighborhood deprivation; each participant's postcode was entered into the relevant and latest UK country databases (England, Scotland, Wales and Northern Ireland) and a quintile rank was determined. Maternal educational level was scored 0 (*college education or below*), or 1 (*university education or above*). Employment status was scored 0 (*no employment*) or 1 (*employment, full or part time*). Neighborhood deprivation was scored 0 (*low quintile*) or 1 (*not deemed low quintile*). For example, the highest score of three indicated living in a low quintile neighborhood, without a paid job, and with educational qualifications below degree level.

*Child Behavioral and Emotional Problems.* The behavioral and emotional problems of the child with IDD were measured using the Strengths and Difficulties Questionnaire (SDQ; Goodman et al., 1997, 1998). The SDQ is a well validated instrument and research with children with IDD and their parents suggests good levels of reliability (Beck et al., 2004a, 2004b; Hastings et al., 2006). Cronbach's alpha coefficient for the total difficulties score was .86 in this present study.

*Maternal resilience.* The Brief Resilience Coping Scale (Sinclair & Wallston, 2004) is designed to assess an adult's ability to recover from stress. The original measure was designed to test resilience in a sample of women with rheumatoid arthritis. The four questions in this scale include: "I actively look for ways to replace losses I encounter in life," "I believe that I can grow in positive ways by dealing with difficult situations," "I look for creative ways to alter difficult situations," "Regardless of what happens to me, I believe I can control my reaction to it." The response scale is a Likert response ranging from 0 (*does not describe*

me at all) to 5 (*describes me very well*). The maximum score is 20, and a higher score indicates greater resilience. According to Sinclair and Wallston (2004) low resilience participants are those who obtain scores lower than 13, while those who scored above 17 are considered highly adaptable. Cronbach's alpha coefficient in the present study was .73 for mothers.

*Maternal Stress.* General parenting stress related to having a child with a disability was measured using a shortened seven-item version of the Parent and Family Problems scale from the Questionnaire on Resources and Stress- short Form (QRSF7: Griffith et al., 2011). Parents were asked to circle either "True" or "False" for each item based on whether the item applied to their family. A total stress score is derived by summing the number of negatively endorsed items (i.e., positively worded items are reverse scored). In the present study, a Kuder-Richardson coefficient for the seven-item scale of .90 was obtained.

*Family Satisfaction Scale.* Family satisfaction was measured by the Family Satisfaction Scale (FSS: Olson & Wilson, 1982), a 14-item scale designed to measure satisfaction on the dimensions of family cohesion and family adaptability (flexibility). Cronbach's alpha coefficient for mothers in this present study, for the total family satisfaction score, including both the cohesion and adaptability subscales was .94.

*Maternal Anxiety and Depression.* Maternal anxiety and depression symptoms over the past seven days were measured by the Hospital Anxiety and Depression Scale (HADS: Zigmond & Snaith, 1983). The HADS has shown good psychometric properties (Hastings et al., 2005) and good levels of reliability when used with mothers of children with IDD (e.g., Hastings & Brown, 2002). In the present sample, Cronbach's alpha coefficients for mothers were .85 for depression and .86 for anxiety.

*Maternal Positive Perceptions.* The Positive Gain Scale (PGS: MacDonald et al., 2010) assesses the positive aspects of raising a child with a disability. The measure consists

of seven items about raising a child with intellectual disability and their putative positive impact on the parent and family. The PGS total score was used in the current study (with lower scores indicating higher levels of positive gain). Cronbach's alpha coefficient for the present sample of mothers was .85.

## Results

To assess maternal resilience as a moderator or as a compensatory factor, multiple regression analyses were conducted for each of the five psychological well-being measures (anxiety, depression, family satisfaction, positive perceptions and parenting/family stress; see Table 3). Relevant demographic variables were selected to be included in each of the five analyses from bivariate analyses (correlations or *t*-tests). Mothers of children with Down's Syndrome reported significantly less stress than other mothers ( $t(310) = 6.49, p < .001$ ), as well as significantly less anxiety ( $t(310) = 4.45, p < .001$ ), and less depression ( $t(310) = 3.89, p < .001$ ), and more family satisfaction ( $t(310) = 2.95, p = .003$ ), than mothers in the study whose child did not have Down's Syndrome. Mothers of children who had a diagnosis of ASD also had higher stress levels ( $t(310) = -4.18, p < .001$ ), higher anxiety ( $t(310) = -4.52, p < .001$ ), higher levels of depression ( $t(310) = -3.72, p < .001$ ), and lower family satisfaction ( $t(310) = 3.16, p = .002$ ), than mothers of children who did not have an ASD diagnosis.

Mothers of male children reported higher stress levels than mothers of female children ( $t(310) = 2.49, p = .01$ ). Mothers of white British ethnicity reported more anxiety ( $t(310) = 2.46, p = .01$ ), and less family satisfaction ( $t(310) = .25, p = .01$ ) than mothers of other ethnicities. Pearson's correlations showed mothers from families with higher SEP reported higher family satisfaction than those mothers from a lower SEP ( $r = .12, p = .03$ ), also families with lower SEP reported lower stress levels ( $r = .14, p = .01$ ), lower anxiety levels ( $r = .11, p = .045$ ), and lower depression levels ( $r = .21, p < .001$ ), than mothers from families

with higher SEP. Older mothers also reported less anxiety ( $r = -.17, p = .002$ ) and less depression ( $r = -.13, p = .03$ ) than younger mothers.

The key predictor variable in each analysis was child behavioral and emotional problems; resilience was entered as a main effect variable, and as an interaction variable with child behavioral and emotional problems. The “PROCESS” custom dialogue box (Hayes, 2012) was installed into SPSS predictive analytics software for the moderated multiple regression analyses. Multicollinearity issues between variables were checked using the Variance Inflation Factor (VIF) and the variables showed no multicollinearity problems (all values  $< 10$ , average  $> 1$ , tolerance  $> 0.1$ ; Bowerman & O’Connell, 1990; Myers, 1990). Predictor variables were automatically mean-centered when using the PROCESS dialogue box (the variable mean is subtracted from every value of the variable).

The moderated multiple regression analysis showed that child behavioral and emotional problems and maternal resilience each had a significant association with maternal well-being outcomes in all five models. There were two potentially relevant interaction terms, one statistically significant (maternal stress) and one close to significance (family satisfaction,  $p = .058$ ) suggesting that the main effects could be interpreted in relation to an interaction effect. Following the recommendation by Aiken and West (1991), a simple slope analysis was conducted to aid interpretation of these two interactions. There was a positive relationship between child behavioral and emotional problems and maternal stress at all three levels of maternal resilience (all  $ps < .001$ ). Thus, the nature of the interaction effect was unclear. Visual inspection of the slopes showed that higher levels of maternal resilience were associated with lower maternal stress when child behavioral and emotional problems were at low levels.

## Discussion

In Study 1, levels of maternal resilience consistently had a significant independent association with maternal positive and negative well-being outcomes when child behavioral and emotional problems were present. Therefore, we found the strongest support for a compensatory model of resilience. Although there were interaction terms (one significant, and one borderline significant) that potentially support a protective factor/moderation model of resilience, the resulting relationships did not follow the predicted pattern of resilience affecting maternal outcomes at high levels of risk (high child behavioral and emotional problems). Instead, we found higher levels of resilience were associated with better maternal outcomes at low levels of child behavioral and emotional problems.

## Study 2

The purpose of the second study was to extend Study 1 by testing the protective and compensatory function of resilience in mothers of children with ASD residing in the USA. First, to test for the protective function of resilience we examined whether maternal resilience moderated the association between child ASD symptom severity and maternal well-being. Next, the direct effect of resilience on maternal well-being was examined using cross-sectional data to test for the compensatory function of maternal resilience and using longitudinal data to test the challenge model of resilience.

## Method

**Participants.** The current sample was from a larger study of parents of children with ASD ( $n = 136$ ). The participants were 99 mothers of children with ASD (84 male) aged between two and 13 years ( $M = 7.78$ ,  $SD = 2.66$ ). The mothers' ages ranged from 25 to 55 years ( $M = 40.00$ ,  $SD = 6.16$ ). Detailed demographic information is shown in Table 4.

## Measures.

*Demographic Questionnaire.* Demographic information was gathered using a questionnaire developed by the research team and included questions about the mother and



their child with ASD. The demographic variables presented in Table 4 were recoded dichotomously: maternal employment was re-coded into two categories (no paid employment vs. employed), ethnicity was coded as white Hispanic or Latino versus all other categories, and education was coded into college level and above versus lower than college level. Other categories were dichotomously coded such as child gender (male vs. female). Three individual indicators were again combined into an index of deprivation. Each indicator was scored dichotomously; educational level was scored 0 (*high school education or below*), or 1 (*vocational education, some college classes, college degree, post college professional degree*). Employment status was scored 0 (*no employment*) or 1 (*employment, full or part time*). Household income was scored as 0 (*low income, to \$24,999*) or 1 (*income above \$24,999*). This is based on the U.S. Department of Health and Human Services, where the poverty guideline is less than \$24,250 for an average of four persons in the household. Total SEP was calculated by summing the scores of these three indicators from the dichotomous coding, a high score indicated low socio economic position.

*Current Child ASD Symptoms.* The severity of child's current ASD symptoms was measured using the parent report version of the Social Responsiveness Scale (SRS: Constantino, Przbeck, Friesen, & Todd, 2000). The SRS is a 65-item scale measuring autistic traits, including social information processing, social use of language, stereotypic/repetitive behaviors/preoccupations, social awareness, and the capacity for reciprocal social response. Responses to the questions are using a four-point Likert scale ranging from 1(*never true*) to 4 (*almost always true*). The responses are summed from the 65-items to produce a total score (index of ASD symptom severity). A higher score indicates a higher level of severity of the child's current ASD symptoms. Cronbach's alpha coefficients were .88 for Time 1 and .87 for Time 2.

1                   *Maternal Resilience.* Resilience was measured using the Ego-Resilience 89 Scale  
2 (ER-89: Block & Kremen, 1996) and is based on their experience with earlier resilience  
3 scales. The ER89 is a 14-item scale focusing on flexibility, curiosity, generosity and social  
4 skills. Sample items include “I quickly get over and recover from being startled” and “I am  
5 more curious than most people.” Participants are asked to respond on a five-point response  
6 scale ranging from 1 (*does not apply at all*) to 4 (*applies very strongly*). Cronbach’s alpha  
7 coefficients for the current study were .71 for Time 1 and .77 for Time 2.  
8  
9

10                   *Depression.* Maternal depressive symptoms were assessed using the Center for  
11 Epidemiological Studies Depression Inventory (CES-D: Devins et al., 1988; Radloff, 1977).  
12 The CES-D is a 20-item self-report questionnaire designed to assess depressive symptoms in  
13 adults. Participants are asked to indicate how frequently they have experienced various  
14 symptoms during the previous week, using a four-point scale 0 (*rarely or none of the time*) to  
15 4 (*most or all of the time*). A high score is indicative of higher depression in mothers; to  
16 achieve this, positively worded items were reverse coded. Previous studies have reported high  
17 internal consistency, adequate test-retest reliability, and good criterion and discriminant  
18 validity (Devins et al., 1988; Radloff, 1977). Cronbach’s alpha coefficients for the current  
19 study were .90 for Time 1 and .91 for Time 2.  
20  
21

22                   *Anxiety.* Maternal anxiety was assessed using the State-Trait Anxiety Inventory  
23 (STAI: Spielberger 1983). The STAI is a 20-item widely used self-report measure of anxiety,  
24 which measures state anxiety (e.g., current temporary experience of anxiety in specific  
25 situations). In the STAI a total score is provided for state anxiety, the score ranges from 20 to  
26 80, and a high score is indicative of higher levels of current anxiety. This measure has been  
27 used previously with parents of children with ASD and has demonstrated good internal  
28 consistency for the STAI (Clifford & Minnes, 2013; Reaven et al., 2015). In this present  
29 study Cronbach’s alpha coefficients for the state scale were .92 for Time 1 and Time 2.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1                   *Family Cohesion.* Family cohesion was assessed using the cohesion subscale of the  
2 Family Adaptability and Cohesion Evaluation Scales IV (FACES: Olsen, 2011). The  
3 cohesion subscale consists of seven items and an example item includes “Family members  
4 consult other family members on personal decisions.” All items are answered using a five-  
5 point scale 1 (*does not describe our family*) to 5 (*very well describes our family*). Higher  
6 scores indicate higher cohesion. The FACES IV scales have demonstrated adequate reliability  
7 and validity (Olsen, 2011). Cronbach’s alpha coefficients were .82 at T1 and T2.  
8  
9

10                   *Benefit Finding.* Benefit finding was measured using the Post Traumatic Growth  
11 Inventory (PTGI: Tedeschi & Calhoun, 1996). The PGTI is a 21-item measure used for  
12 assessing positive outcomes when “traumatic events” have been experienced. Participants  
13 responded to each item using a five-point Likert scale ranging from 0 (*I did not experience*  
14 *this change as a result of the incident*) to 5 (*I experienced this change to a very great degree*  
15 *as a result of this incident*). In this study, the “incident” is referring to having a child with  
16 ASD. The PGTI has good reliability and validity. Cronbach’s alpha coefficients for the full  
17 scale in this study were .89 for Time 1 and .89 for Time 2.  
18  
19

20                   *Loneliness.* Maternal loneliness was measured using the revised version of the UCLA  
21 Loneliness Scale (Peplau, & Cutrona, 1980). The 20-item scale asks participants to describe  
22 their feelings of loneliness through non-lonely items (e.g., *I feel part of a group of friends* and  
23 *I lack companionship*). The scale has been shown to have good reliability (Russell, 1996). A  
24 higher score reflects a higher level of loneliness reported by the mother. Cronbach’s alpha  
25 coefficients for this current study were .93 for Time 1 and Time 2.  
26  
27

28                   **Procedure.** Participants were recruited through online resources throughout the  
29 United States (e.g., blogs, Facebook groups, online autism support groups) and through word  
30 of mouth. Mothers who expressed interest in the study were provided with further details  
31 about the study and were emailed a unique link to complete all of the questionnaires online.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

After clicking the link, participants first read and electronically signed the informed consent and then proceeded to complete questionnaires. Upon completion of the survey, parents were mailed a \$10 gift card for a national retailer. Approximately six months later, participants were contacted to complete the second part of the survey. Once again, they were emailed a unique link to the survey and completed the questionnaires online. After completion, parents were mailed another \$10 gift card for a national retailer. Due to the nature of the recruitment methods, we are unable to determine the overall response rate for this survey.

## Results

To assess maternal resilience in cross sectional and longitudinal data, multiple regression analyses were conducted for each of the psychological well-being measures (depression, family cohesion, anxiety, loneliness, benefit finding). Demographic variables that were statistically significant with outcome variables were selected to be included in each of the analyses from bivariate analyses (correlations or *t*-tests). Mothers who had no additional children with ASD in the house had higher scores on the FACES cohesion scale ( $t(89) = 2.15, p = .03$ ). Therefore, having additional children with ASD in the house was included as a control variable when family cohesion served as the outcome variable.

All of the continuous variables were examined for normality using Kolmogorov-Smirnov tests; this showed that all variables were normally distributed and suitable for parametric analysis. Multicollinearity issues between variables were checked using the Variance Inflation Factor (VIF) and the variables showed no multicollinearity problems (all values  $<10$ , average  $>1$ , tolerance  $>0.1$ ) (Bowerman & O'Connell, 1990; Myers, 1990).

*Cross-sectional analyses.* Cross-sectional analyses were conducted to establish whether maternal resilience functioned as a moderator between the severity of the child's current ASD symptoms and maternal outcomes (see Table 5). Using Time 1 data, moderated multiple regression analyses for each of the maternal well-being outcome variables were

1 conducted. The key predictor entered in each analysis was the severity of the child's current  
2 ASD symptoms. Maternal resilience was entered in the analyses both as a main effect and as  
3 an interaction variable. Significant demographic variables were entered as control variables,  
4 when appropriate. The "PROCESS" custom dialogue box (Hayes, 2012) was installed into  
5 SPSS predictive analytics software for the moderated multiple regression analyses. Predictor  
6 variables were automatically mean-centered when using the PROCESS dialogue box (the  
7 variable mean is subtracted from every value of the variable).  
8  
9  
10  
11  
12  
13  
14  
15

16  
17 In the cross-sectional analyses, maternal resilience did not moderate outcomes in any  
18 of the models. However, maternal resilience had a significant independent association with  
19 maternal depression, anxiety, loneliness, and family cohesion. The severity of the child's  
20 current ASD symptoms, as reported by the mother, was a significant independent predictor  
21 for the maternal outcomes of depressive symptoms, anxiety, and loneliness.  
22  
23  
24  
25  
26  
27  
28

29 *Longitudinal analyses.* Longitudinal analyses were conducted to identify if maternal  
30 resilience at Time 1 predicted later maternal well-being at Time 2 (see Table 6). These  
31 longitudinal analyses again used regression analyses for each of the maternal well-being  
32 outcome variables. Time 2 well-being outcomes were entered as criterion variables in the  
33 regression analyses. The key predictor variables entered in each analysis were Time 1  
34 maternal resilience, Time 1 severity of child's current ASD symptoms, the Time 1 score for  
35 the criterion variables, and the relevant demographic variable selected from the cross-  
36 sectional analyses.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 The longitudinal multiple regression analyses showed that maternal resilience did not  
49 act as a significant predictor of maternal well-being outcomes at Time 2. The only variable to  
50 make an independent contribution to the prediction of later maternal well-being was the  
51 severity of the child's current ASD symptoms: mothers reported lower benefit finding scores  
52 at Time 2 when their child with ASD had more severe symptoms at Time 1.  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Discussion

The cross-sectional analyses in Study 2 led to similar findings to those from Study 1: maternal resilience had a significant independent association with two maternal outcome variables including depression and family cohesion. Therefore, we again found the strongest support for a compensatory model of resilience. The longitudinal analyses found maternal resilience did not act as a significant predictor of maternal well-being outcomes over time.

In the cross-sectional analysis, the severity of the child's current ASD symptoms, as reported by the mother, had a significant independent association with both anxiety and loneliness in mothers of children with ASD. This is consistent with previous research which found that severity of the child's ASD symptoms had a negative effect on maternal outcomes, such as anxiety, (e.g., Firth & Dryer, 2013). The longitudinal analyses also showed high severity of the child's ASD symptoms predicted lower benefit-finding in mothers over time.

## General Discussion

The purpose of the present research was to examine whether resilience, the ability to resist the negative impact of adverse situations, promotes psychological well-being in mothers of children with DD, including ASD. Study 1 found support for a compensatory model of resilience whereby maternal resilience was associated with better psychological well-being for mothers of children with DD. Contrary to predictions, we found limited support for the protective model of resilience. Higher levels of resilience only served as a protective factor in the context of low levels of child behavioral and emotional problems and reliably in only one of the regression models tested. Study 2 explored the compensatory and protective models of resilience in a sample of mothers of children with ASD. Similar to Study 1, support for the compensatory model of resilience was found. Using longitudinal data over six months in Study 2, no support was found for a temporal relationship between resilience and later maternal psychological outcomes.

1           The compensatory model of resilience posits that resilience has a direct impact on  
2 well-being outcomes and we found support for this model in both studies. In Study 1,  
3 maternal resilience was associated with all study outcomes in the expected direction. That is,  
4 greater levels of resilience were associated with better well-being in mothers of children with  
5 IDD. In Study 2, using a different measure of resilience, higher levels of resilience was  
6 associated with less depressive symptoms, lower anxiety, less loneliness, and better family  
7 functioning in mothers of children with ASD. These findings support a growing body of  
8 literature showing that positive characteristics (e.g., optimism, hope) of a parent of child with  
9 DD are associated with fewer negative outcomes and increased positive outcomes (Ekas,  
10 Pruitt, & McKay, 2016; Ekas, Keyton, Pruitt, Ghilain, & Alessandri, 2015; Ekas,  
11 Lickenbrock, & Whitman, 2010; Hastings & Taunt, 2002; Lloyd & Hastings, 2009). One  
12 aspect that can be explored further is to examine the attributes of “resilient mothers.” For  
13 example, there may be some overlap between the construct of resilience, as measured in this  
14 research, and other constructs commonly associated with maternal well-being. For example,  
15 one question on the resilience measure (ER-89) asks “I usually succeed in making a favorable  
16 impression,” which may be linked to self-esteem.

17           Studies 1 and 2 also examined the protective model of resilience wherein resilience is  
18 hypothesized to “buffer” (i.e., moderate) the negative effects of child behavioral and  
19 emotional problems or child ASD symptom severity. In Study 1, although we found a  
20 significant interaction between maternal resilience and child behavioral and emotional  
21 problems predicting maternal stress, the direction of effects was contrary to our predictions.  
22 We found higher levels of resilience were associated with better maternal outcomes at low  
23 levels of child behavioral and emotional problems when we expected resilience to be  
24 protective at higher levels of child behavioral and emotional problems. In Study 2, we did not  
25 find any evidence for the protective model in mothers of children with ASD. The unexpected

findings in relation to the regression models in Study 1 may relate to the third theoretical perspective on resilience introduced earlier - the challenge model. It is possible that in a sample of mothers of children with DD, “low risk child behavioral problems” is the equivalent of “high risk child behavioral problems” when compared to families of children without DD. Therefore, high risk in the current study might constitute “extremely high risk” (having a child with IDD and with significant levels of behavior problems). Ongoing exposure to risk in these families may have increased their resilience over time and this may be reflected in the data from this cross-sectional study. However, at this point the families may have already built up significant resilience at a high or extremely high risk in terms of their child’s behavior. We also tested if the compensatory model held over time, addressing the key causality question of temporal precedence. We found no evidence for a compensatory effect over time in Study 2.

### **Limitations and Future Research**

There are several limitations that warrant discussion. First, upon examining the resilience measures available it was clear there is no resilience measure available that clearly pinpointed the concept of resilience from the definitions and evidence available. However, this is not a direct limitation unique to our study design but comes from a much broader issue of the difficulty in defining and measuring resilience. The resilience measure selected in Study 1 was a short, four question measure which may have reduced the internal consistency; although the internal consistency was still acceptable (.73) within statistical recommendations (Hayes, 2012). The definitions of resilience in previous literature include the idea of bouncing back “the ability to withstand hardship and rebound from adversity, becoming more strengthened and resourceful” (Walsh, 1998, 2006, p. 263). The questions addressed in the measure of resilience seem to address factors associated with resilient outcomes, such as handling stress in an adaptive manner. The resilience measure for this study was selected as it



1 best reflected the core concept of resilience and past definitions, it also scored highly in a  
2 reliability and validity assessment (Windle, 2011). Study 2 was the first to use the ER-89  
3 with mothers of children with ASD, and therefore this ER-89 measure may not be capturing  
4 resilience in these families. However, Cronbach's alpha coefficients were good in the current  
5 study for the ER-89 (.71 for Time 1 and .77 for Time 2). More research is needed to clarify  
6 what these measures of resilience are capturing. The attributes of "resilience" captured in the  
7 ER-89 suggests these do have an association with maternal outcomes. For example,  
8 identifying one's ability to "get over things quickly" suggests successful coping mechanisms.  
9 Another consideration is the theory and discussion in literature of whether resilience can be  
10 captured in a measure, or if it is a process that unfolds over time. Social psychologists  
11 generally think of resilience as a trait-like phenomenon whereas developmental psychologists  
12 tend to think of resilience as something that is demonstrated over time (e.g., see Masten,  
13 Powell & Luthar, 2003; Masten & Obradović, 2006).

14 Another limitation is that mothers provided all the data in this research which means  
15 there was a lack of source variance. To address this, future research will need to incorporate  
16 independent or multiple informant approaches for key constructs (e.g., child behavioral and  
17 emotional problems).

18 Finally, Study 1 was limited by its cross-sectional design. This is problematic since  
19 temporal precedence has not been established. For example, it may be the case that mothers  
20 with higher levels of psychological distress become less resilient. Study 2 attempted to  
21 address this concern by including longitudinal data. However, the findings were non-  
22 significant. One possible explanation for the non-significant findings in the longitudinal  
23 analyses, is the time period between the two data points. The six-month time period between  
24 the two data points may not have been enough time for well-being outcomes to change.  
25 Previous research with families of children with ASD (e.g., Shattuck et al., 2007) conducted

longitudinal analyses for a time period of four and a half years, to reflect the changes seen in child behavioral and emotional problems over adolescence. This study showed that over time as child behavioral and emotional problems decreased, maternal well-being improved. Therefore, future research should consider a longer time frame for follow up, to follow changes to child behavioral problems, and also to see if resilience has built in this time, and if these predict maternal well-being.

### **Clinical Implications**

Despite these limitations, the results of two studies provide initial support for the compensatory model of resilience in mothers of children with DD, including ASD. The finding that maternal resilience is associated with maternal well-being may be used to inform clinical practice in improving well-being in mothers of children with DD, including ASD. In particular, it would be beneficial to directly try to build resilience in mothers to improve their well-being. As one aspect of resilience addresses mothers' positivity, this could potentially be the appropriate target for interventions. A meta-analysis showed that a number of interventions have been successful in the general population in improving positivity, and therefore well-being (Sin & Lyubomirsky, 2009). Several positive psychology interventions were found to be effective in improving well-being, such as a person identifying their strengths and using their signature strengths in new ways (Seligman, Steen, Park, & Peterson, 2005); cognitive strategies for example replaying positive experiences and self-monitoring well-being (Fava, Rafanelli, Cazzaro, Conti, & Grandi, 1998); and practicing emotional skills such as mindfulness and acceptance (Bedard et al., 2003; Grossman, Tiefenthaler-Gilmer, Raysz, & Kesper, 2007). It may be that the previous research into what we suspect to be aspects of resilience, such as positivity, may help form a resilience intervention, which would also help develop a clear concept of resilience. Exploration of these types of interventions

which focus on aspects of resilience for families of children with DD is needed in future research.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Table 1

*Study 1: Mothers' Demographic Information*

| Variable                      |  | <i>n</i> | <i>Percent</i> |
|-------------------------------|--|----------|----------------|
| Postcode deprivation quintile | 1 – Least Deprived   | 68       | 22 %           |
|                               | 2  | 48       | 15 %           |
|                               | 3  | 42       | 14 %           |
|                               | 4  | 56       | 18 %           |
|                               | 5- Most Deprived   | 69       | 22 %           |
| Education level               | No formal educational qualification                                | 11       | 4 %            |
|                               | Fewer than 5 GCSE's/ or levels or equivalent                       | 21       | 7 %            |
|                               | 3 or more a levels (NVQ 3) or equivalent                           | 43       | 14 %           |
|                               | University degree  | 124      | 40 %           |
|                               | Masters or doctoral degree   | 51       | 16 %           |
| Employment Status             | No paid employment   | 125      | 40 %           |
|                               | Part time  | 125      | 40 %           |
|                               | Full time  | 32       | 10 %           |
|                               | Self -employed (full/part time)                                    | 30       | 10 %           |
| Ethnicity                     | White British  | 262      | 84 %           |
|                               | White Irish  | 5        | 2 %            |
|                               | White Welsh  | 24       | 8 %            |
|                               | Other White background   | 10       | 3 %            |
|                               | Mixed White and Asian  | 2        | 1 %            |
|                               | Other Mixed background   | 3        | 1 %            |
|                               | Black/ Black British- Caribbean                                    | 1        | <1 %           |
|                               | White and Black Caribbean  | 1        | <1 %           |
|                               | Asian/Asian British (Bangladeshi),<br>Asian/Asian British (Indian) | 1        | <1 %           |
|                               | Other Asian background   | 2        | 1 %            |

Table 2

*Study 1: Children's Demographic Information*

| Variable   | <i>n</i> | <i>Percent</i> |
|--|----------|----------------|
| Male   | 227      | 73 %           |
| Autism diagnosis   | 171      | 55 %           |
| Various diagnoses and causes of their DD*  | 93       | 30 %           |
| Down's Syndrome  | 48       | 15 %           |
| Disability from birth rather than acquired   | 263      | 84 %           |
| Additional health condition  | 162      | 52 %           |
| Secondary diagnosis given  | 123      | 39 %           |
| Can feed themselves/ feed themselves with help   | 295      | 95 %           |
| Can dress themselves or dress with help  | 259      | 83 %           |
| Can walk upstairs without help/ by themselves  | 256      | 82 %           |
| Can wash themselves or wash with help  | 252      | 81 %           |
| Hearing impairment or deaf   | 55       | 18 %           |
| Children did not use speech  | 39       | 13 %           |
| *Examples include; no specific diagnosis such as SWAN- Syndrome Without A Name, genetic syndromes such as Fragile X Syndrome, and other diagnoses such as Global Developmental Delay |          |                |

Table 3

*Study 1: Moderated Multiple Regression Analyses Models for the Five Maternal Psychological Well-being Measures*

| <i>n</i> = 312  | <u>Maternal Stress</u><br><i>R</i> <sup>2</sup> = .37 |                 | <u>Family Satisfaction</u><br><i>R</i> <sup>2</sup> = .29 |                 | <u>Maternal Anxiety</u><br><i>R</i> <sup>2</sup> = .25 |                 | <u>Maternal Depression</u><br><i>R</i> <sup>2</sup> = .28 |                 | <u>Positive Perceptions</u><br><i>R</i> <sup>2</sup> = .16 |                 |
|---|---|-----------------|---|-----------------|--|-----------------|---|-----------------|--|-----------------|
| <u>Variable</u>   | <i>β</i>  | <i>p</i>        | <i>β</i>  | <i>p</i>        | <i>β</i>   | <i>p</i>        | <i>β</i>  | <i>p</i>        | <i>β</i>   | <i>p</i>        |
| Age of mother   | .01   | .43             | -.20  | .004            | -.07   | .06             | -.03  | .43             | .09  | <b>.04</b>      |
| Autism present  | -.31  | .21             | -.65  | .53             | .45  | .48             | .24   | .69             | .18  | .87             |
| Down's Syndrome present   | -.82  | <b>.03</b>      | .76   | .62             | -.68   | .35             | -.54  | .46             | .59  | .56             |
| Gender of child   | -.45  | .06             | -.01  | .99             | -.52   | .33             | .25   | .64             | .04  | .96             |
| SEP   | .25   | .07             | -1.19   | <b>.04</b>      | .37  | .25             | .95   | <b>.003</b>     | .33  | .39             |
| White ethnicity   | .37   | .26             | -2.42   | <b>.03</b>      | 1.51   | <b>.03</b>      | .42   | .54             | .44  | .53             |
| Child behavior problems (centered)  | .13   | <b>&lt;.001</b> | -.29  | <b>&lt;.001</b> | .17  | <b>&lt;.001</b> | .16   | <b>&lt;.001</b> | .09  | <b>.02</b>      |
| Maternal resilience (centered)  | -.08  | <b>.004</b>     | .85   | <b>&lt;.001</b> | -.27   | <b>&lt;.001</b> | -.39  | <b>&lt;.001</b> | -.41   | <b>&lt;.001</b> |
| Maternal resilience x Child behavioral and emotional problems (interaction) | .01   | <b>.04</b>      | -.04  | .06             | .01  | .62             | .00   | .77             | -.01   | .30             |

*Note: Significant (*p* < .05) associations between variables are in boldface.*

Table 4

*Study 2: Mothers' Demographic Information*

| Variable          |  | <i>n</i> | <i>Percent</i> |
|-------------------|--|----------|----------------|
| Household income  | \$24,999 or less (low income)                              | 10       | 10 %           |
|                   | \$25,000 and above (mean or high income)                   | 90       | 90 %           |
| Education level   | High school (grades 10-12)                                 | 17       | 17 %           |
|                   | Vocational education or some college classes               | 21       | 21 %           |
|                   | College degree   | 41       | 41 %           |
|                   | Post college professional degree (MA, PhD, MD, Law, other) | 16       | 16 %           |
| Employment Status | No paid employment   | 45       | 46 %           |
|                   | Part time  | 20       | 20 %           |
|                   | Full time  | 34       | 34 %           |
| Ethnicity         | White  | 86       | 87 %           |
|                   | Hispanic or Latino   | 12       | 12 %           |
|                   | Hispanic and/ or Latin American                            | 10       | 10 %           |
|                   | Black and/ or African American                             | 2        | 2 %            |
|                   | Native American or Aleutian Islander/ Eskimo               | 1        | 1 %            |
|                   | Asian or Pacific Islander                                  | 1        | 1 %            |
| Marital Status    | Other  | 1        | 1 %            |
|                   | Single/ separated/ widowed/ divorced                       | 16       | 17 %           |
|                   | Married  | 83       | 83 %           |

Table 5  
 Study 2: Time 1 Cross Sectional Analysis of Maternal Well-being Outcomes

| <i>Time 1 Maternal well-being outcomes</i>             | <u>Depression</u><br>R <sup>2</sup> = .15<br>n = 92 |            | <u>Anxiety</u><br>R <sup>2</sup> = .14<br>n = 92 |             | <u>Cohesion</u><br>R <sup>2</sup> = .18<br>n = 88 |             | <u>Benefit finding</u><br>R <sup>2</sup> = .03<br>n = 91 |          | <u>Loneliness</u><br>R <sup>2</sup> = .15<br>n = 91 |             |
|--|---|------------|--|-------------|---|-------------|--|----------|---|-------------|
| <u>Variable</u>  | $\beta$   | <i>p</i>   | $\beta$  | <i>P</i>    | $\beta$   | <i>p</i>    | $\beta$  | <i>p</i> | $\beta$   | <i>p</i>    |
| Additional child with ASD                              | --  | --         | --   | --          | -2.65   | .06         | --   | --       | --  | --          |
| Severity of ASD current symptoms                       | <b>.11</b>  | <b>.01</b> | <b>.117</b>                                      | <b>.01</b>  | .00   | .96         | .07  | .12      | <b>.13</b>  | <b>.004</b> |
| Maternal resilience                                    | <b>-.45</b>   | <b>.03</b> | <b>-.445</b>                                     | <b>.047</b> | <b>.34</b>  | <b>.002</b> | .15  | .54      | <b>-.48</b>   | <b>.04</b>  |
| Severity of ASD current symptoms x maternal resilience | .00   | .96        | .001   | .86         | .00   | .53         | -.00   | .76      | .01   | .30         |

*Note. Significant ( $p < .05$ ) associations between variables are in boldface*



Table 6

*Study 2: Longitudinal Analysis of Maternal Well-being Outcomes*

| <i>Time 2 Maternal well-being outcomes</i>    | <u>Depression</u> |                 | <u>Anxiety</u> |                 | <u>Cohesion</u> |                 | <u>Benefit finding</u> |                 | <u>Loneliness</u> |                 |
|---|-------------------|-----------------|----------------|-----------------|-----------------|-----------------|------------------------|-----------------|-------------------|-----------------|
|   | $R^2 = .36$       |                 | $R^2 = .23$    |                 | $R^2 = .59$     |                 | $R^2 = .42$            |                 | $R^2 = .65$       |                 |
|   | $n = 70$          |                 | $n = 70$       |                 | $n = 66$        |                 | $n = 69$               |                 | $n = 68$          |                 |
| <u>Variable</u>                               | $\beta$           | $p$             | $\beta$        | $p$             | $\beta$         | $p$             | $\beta$                | $p$             | $\beta$           | $p$             |
| Additional child with ASD                     | --                | --              | --             | --              | .94             | .43             | --                     | --              | --                | --              |
| Maternal resilience Time 1                    | -.27              | .20             | -.23           | .32             | .15             | .10             | .09                    | .70             | -.14              | .41             |
| Severity of child ASD current symptoms Time 1 | .05               | .18             | .03            | .48             | -.00            | .85             | <b>-.09</b>            | <b>.046</b>     | .04               | .21             |
| Outcome Time 1                                | <b>.46</b>        | <b>&lt;.001</b> | .45            | <b>&lt;.001</b> | <b>.73</b>      | <b>&lt;.001</b> | <b>.69</b>             | <b>&lt;.001</b> | <b>.77</b>        | <b>&lt;.001</b> |

Note. Significant ( $p < .05$ ) associations between variables are in boldface, longitudinal results are shown after accounting the Time 1 score in each outcome

### References

- Aiken, L. S., & West, S. G. (1991). *Multiple regression: Testing and interpreting interactions*. Sage Publications, Inc.
- Andrews, G., Page, A. C., & Neilson, M. (1993). Sending your teenagers away: controlled stress decreases neurotic vulnerability. *Archives of General Psychiatry*, 50(7), 585-589.
- Baker, B. L., McIntyre, L. L., Blacher, J., Crnic, K., Edelbrock, C., & Low, C. (2003). Pre-school children with and without developmental delay: behaviour problems and parenting stress over time. *Journal of Intellectual Disability Research*, 47(4-5), 217-230.
- Bayat, M. (2007). Evidence of resilience in families of children with autism. *Journal of Intellectual Disability Research*, 51, 702-714.
- Beck, A., Daley, D., Hastings, R. P., & Stevenson, J. (2004). Mothers' expressed emotion towards children with and without intellectual disabilities. *Journal of Intellectual Disability Research*, 48(7), 628-638.
- Beck, A., Hastings, R. P., Daley, D., & Stevenson, J. (2004). Pro-social behaviour and behaviour problems independently predict maternal stress. *Journal of Intellectual and Developmental Disability*, 29(4), 339-349.
- Bedard, M., Felteau, M., Mazmanian, D., Fedyk, K., Klein, R., Richardson, J., Parkinson, W., & Minthorn-Biggs, M. B. (2003). Pilot evaluation of a mindfulness-based intervention to improve quality of life among individuals who sustained traumatic brain injuries. *Disability and Rehabilitation*, 25(13), 722-731.
- Blacher, J., & McIntyre, L. L. (2006). Syndrome specificity and behavioural disorders in

young adults with intellectual disability: Cultural differences in family impact.

*Journal of Intellectual Disability Research*, 50(3), 184-198.

Block, J., & Kremen, A. M. (1996). IQ and ego-resiliency: conceptual and empirical connections and separateness. *Journal of Personality and Social Psychology*, 70(2), 349-361.

Bowerman, B. L., & O'Connell, R. T. (1990). *Linear statistical models: An applied approach* (2<sup>nd</sup> ed.). Belmont, CA: Duxbury.

Brook, J. S., Whiteman, M., Gordon, A. S., & Cohen, P. (1986). Dynamics of childhood and adolescent personality traits and adolescent drug use. *Developmental Psychology*, 22(3), 403-414.

Brook, J. S., Whiteman, M., Gordon, A. S., & Cohen, P. (1989). Changes in drug involvement: A longitudinal study of childhood and adolescent determinants. *Psychological Reports*, 65(3), 707-726.

Clifford, T., & Minnes, P. (2013). Who participates in support groups for parents of children with autism spectrum disorders? The role of beliefs and coping style. *Journal of Autism and Developmental Disorders*, 43(1), 179-187.

Constantino, J. N., Przybeck, T., Friesen, D., & Todd, R. D. (2000). Reciprocal social behavior in children with and without pervasive developmental disorders. *Journal of Developmental & Behavioral Pediatrics*, 21(1), 2-11.

Dabrowska, A., & Pisula, E. (2010). Parenting stress and coping styles in mothers and fathers of pre-school children with autism and Down syndrome. *Journal of Intellectual Disability Research*, 54(3), 266-280.

- Devins, G. M., Orme, C. M., Costello, C. G., Binik, Y. M., Frizzell, B., Stam, H. J., & Pullin, W. M. (1988). Measuring depressive symptoms in illness populations: Psychometric properties of the Center for Epidemiologic Studies Depression (CES-D) scale. *Psychology and Health*, 2(2), 139-156.
- Eisenhower, A. S., Baker, B. L., & Blacher, J. (2005). Preschool children with intellectual disability: syndrome specificity, behaviour problems, and maternal well-being. *Journal of Intellectual Disability Research*, 49(9), 657-671.
- Ekas, N.V., Keylon, L., Pruitt, M., Ghilain, C., & Alessandri, M. (2015). The power of positivity: Predictors of relationship satisfaction for parents of children with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 45, 1997-2007.
- Ekas, N.V., Lickenbrock, D.M., & Whitman, T.L. (2010). Optimism, social support, and well-being among mothers of children with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 40, 1274-1284.
- Ekas, N.V., Pruitt, M.M., & McKay, E. (2016). Hope, social relations, and depression symptoms in mothers of children with autism spectrum disorder. *Research in Autism Spectrum Disorders*, 29-30, 8-18.
- Ello, L. M., & Donovan, S. J. (2005). Assessment of the relationship between parenting stress and a child's ability to functionally communicate. *Research on Social Work Practice*, 15(6), 531-544.
- Fava, G. A., Rafanelli, C., Cazzaro, M., Conti, S., & Grandi, S. (1998). Well-being therapy. A novel psychotherapeutic approach for residual symptoms of affective disorders. *Psychological Medicine*, 28(02), 475-480.
- Fergus, S., & Zimmerman, M. A. (2005). Adolescent resilience: A framework for

1 understanding healthy development in the face of risk. *Annual Review of Public*  
2 *Health*, 26, 399-419.  
3

4  
5 Firth, I., & Dryer, R. (2013). The predictors of distress in parents of children with autism  
6 spectrum disorder. *Journal of Intellectual and Developmental Disability*, 38(2), 163-  
7 171.  
8  
9

10  
11 Fletcher, D., & Sarkar, M. (2013). Psychological resilience. *European Psychologist*, 18, 12-  
12 23.  
13

14 Friborg, O., Hjemdal, O., Rosenvinge, J. H., & Martinussen, M. (2003). A new rating scale  
15 for adult resilience: what are the central protective resources behind healthy  
16 adjustment? *International Journal of Methods in Psychiatric Research*, 12(2), 65-76.  
17  
18

19 Gerstein, E. D., Crnic, K., Blacher, J., & Baker, B. L. (2009). Resilience and the course of  
20 daily parenting stress in families of young children with intellectual disabilities.  
21 *Journal of Intellectual Disability Research*, 53(12), 981-997.  
22  
23  
24

25 Goodman, R. (1997). The Strengths and Difficulties Questionnaire: A research note. *Journal*  
26 *of Child Psychology and Psychiatry*, 38(5), 581-586.  
27  
28

29 Goodman, R., Meltzer, H., & Bailey, V. (1998). The Strengths and Difficulties  
30 Questionnaire: A pilot study on the validity of the self-report version. *European Child*  
31 *and Adolescent Psychiatry*, 7(3), 125-130.  
32  
33

34 Griffith, G.M., Hastings, R.P., Oliver, C., Howlin, P., Moss, J., Petty, J., & Tunnicliffe, P.  
35 (2011). Psychological wellbeing in parents of children with Angelman, Cornelia de  
36 Lange and Cri du Chat syndromes. *Journal of Intellectual Disability Research*, 55(4),  
37 397-410.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- Grossman, P., Tiefenthaler-Gilmer, U., Raysz, A., & Kesper, U. (2007). Mindfulness training as an intervention for fibromyalgia: evidence of postintervention and 3-year follow-up benefits in well-being. *Psychotherapy and Psychosomatics*, 76(4), 226-233.
- Gutierrez, L. M. (1994). Beyond coping: An empowerment perspective on stressful life events. *Journal of Sociology and Social Welfare*, 21, 201-220.
- Hastings, R. P. (2016). Do children with intellectual and developmental disabilities have a negative impact on other family members? The case for rejecting a negative narrative. *International Review of Research in Developmental Disabilities*, 50, 165-194.
- Hastings, R. P., & Brown, T. (2002). Behavior problems of children with autism, parental self-efficacy, and mental health. *American Journal on Mental Retardation*, 107, 222-232.
- Hastings, R. P., & Johnson, E. (2001). Stress in UK families conducting intensive home-based behavioral intervention for their young child with autism. *Journal of Autism and Developmental Disorders*, 31(3), 327-336.
- Hastings, R. P., & Taunt, H. M. (2002). Positive perceptions in families of children with developmental disabilities. *American Journal on Mental Retardation*, 107, 116-127.
- Hastings, R. P., Daley, D., Burns, C., & Beck, A. (2006). Maternal distress and expressed emotion: Cross-sectional and longitudinal relationships with behavior problems of children with intellectual disabilities. *American Journal on Mental Retardation*, 111, 48-61.
- Hastings, R. P., Kovshoff, H., Brown, T., Ward, N. J., Degli Espinosa, F., & Remington, B. (2005). Coping strategies in mothers and fathers of preschool and school-age children with autism. *Autism*, 9(4), 377-391.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Hayes, A. F. (2012). PROCESS: A versatile computational tool for observed variable mediation, moderation, and conditional process modelling [White paper]. Retrieved from <http://www.afhayes.com/public/process2012.pdf>
- Hayes, S. A., & Watson, S. L. (2013). The impact of parenting stress: A meta-analysis of studies comparing the experience of parenting stress in parents of children with and without autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 43(3), 629-642.
- Herring, S., Gray, K., Taffe, J., Tonge, B., Sweeney, D., & Einfeld, S. (2006). Behaviour and emotional problems in toddlers with pervasive developmental disorders and developmental delay: Associations with parental mental health and family functioning. *Journal of Intellectual Disability Research*, 50(12), 874-882.
- Hill-Chapman, C. R., Herzog, T. K., & Maduro, R. S. (2013). Aligning over the child: Parenting alliance mediates the association of autism spectrum disorder atypicality with parenting stress. *Research in Developmental Disabilities*, 34(5), 1498-1504.
- Lecavalier, L., Leone, S., & Wiltz, J. (2006). The impact of behaviour problems on caregiver stress in young people with autism spectrum disorders. *Journal of Intellectual Disability Research*, 50(3), 172-183.
- Lloyd, T., & Hastings, R. P. (2009). Hope as a psychological resilience factor in mothers and fathers of children with intellectual disabilities. *Journal of Intellectual Disability Research*, 53, 957-968.
- Lyons, D. M., & Parker, K. J. (2007). Stress inoculation-induced indications of resilience in monkeys. *Journal of Traumatic Stress*, 20(4), 423-433.

- MacDonald, E. E., Hastings, R. P., & Fitzsimons, E. (2010). Psychological acceptance mediates the impact of the behaviour problems of children with intellectual disability on fathers' psychological adjustment. *Journal of Applied Research in Intellectual Disabilities*, 23(1), 27-37.
- Masten, A. S., & Obradović, J. (2006). Competence and resilience in development. *Annals of the New York Academy of Sciences*, 1094(1), 13-27
- Masten, A. S., Powell, J. L., & Luthar, S. S. (2003). A resilience framework for research, policy, and practice. *Resilience and vulnerability: Adaptation in the context of childhood adversities*, 1-25.
- McConnell, D., & Savage, A. (2015). Stress and Resilience Among Families Caring for Children with Intellectual Disability: Expanding the Research Agenda. *Current Developmental Disorders Reports*, 2(2), 100-109.
- Myers, R. (1990). *Classical and Modern Regression With Applications* (2<sup>nd</sup> ed.). Boston, MA: Duxbury.
- Oldehinkel, A. J., & Ormel, J. (2014). A longitudinal perspective on childhood adversities and onset risk of various psychiatric disorders. *European Child and Adolescent Psychiatry*, 24, 641-650.
- Olson, D. (2011). FACES IV and the circumplex model: Validation study. *Journal of Marital and Family Therapy*, 37(1), 64-80.
- Olson, D.H., Wilson, M. (1982) *Family satisfaction*. In: Olson, D.H., McCubbin, H.I., Barnes, H., Larsen, A., Muxen, M., Wilson, M., editors. *Family inventories: Inventories used in a national survey of families across the family life cycle*. Family



social science. St. Paul (MN): University of Minnesota.

Pakenham, K. I., Samios, C., & Sofronoff, K. (2005). Adjustment in mothers of children with Asperger syndrome An application of the double ABCX model of family adjustment. *Autism, 9*(2), 191-212.

Peer, J. W., & Hillman, S. B. (2014). Stress and resilience for parents of children with intellectual and developmental disabilities: a review of key factors and recommendations for practitioners. *Journal of Policy and Practice in Intellectual Disabilities, 11*(2), 92-98.

Peplau, L. A., & Cutrona, C. E. (1980). The revised UCLA Loneliness Scale: Concurrent and discriminant validity evidence. *Journal of Personality and Social Psychology, 39*(3), 472-480.

Radloff, L. S. (1977). The CES-D scale a self-report depression scale for research in the general population. *Applied Psychological Measurement, 1*(3), 385-401.

Reaven, J., Washington, L., Moody, E. J., Stern, J. A., Hepburn, S. L., & Blakeley-Smith, A. (2015). Examining the Relationship Between Parental Anxiety and Treatment Response in Children and Adolescents with Autism Spectrum Disorder and Anxiety. *Journal of Autism and Developmental Disorders, 45*(8), 2464-2473.

Ruiz-Robledillo, N., De Andrés-García, S., Pérez-Blasco, J., González-Bono, E., & Moya-Albiol, L. (2014). Highly resilient coping entails better perceived health, high social support and low morning cortisol levels in parents of children with autism spectrum disorder. *Research in Developmental Disabilities, 35*(3), 686-695.

Rutter, M. (1987). Psychosocial resilience and protective mechanisms. *American Journal of*

*Orthopsychiatry*, 57(3), 316-331.

Rutter, M. (2006). Implications of resilience concepts for scientific understanding. *Annals of the New York Academy of Sciences*, 1094(1), 1-12.

Scorgie, K., & Sobsey, D. (2000). Transformational outcomes associated with parenting children who have disabilities. *Mental Retardation*, 38(3), 195-206.

Seery, M. D., Holman, E. A., & Silver, R. C. (2010). Whatever does not kill us: cumulative lifetime adversity, vulnerability, and resilience. *Journal of Personality and Social Psychology*, 99(6), 1025-1041.

Seligman, M. E., Steen, T. A., Park, N., & Peterson, C. (2005). Positive psychology progress: empirical validation of interventions. *American Psychologist*, 60(5), 410-421.

Shattuck, P. T., Seltzer, M. M., Greenberg, J. S., Orsmond, G. I., Bolt, D., Kring, S., ... & Lord, C. (2007). Change in autism symptoms and maladaptive behaviors in adolescents and adults with an autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 37(9), 1735-1747

Sin, N. L., & Lyubomirsky, S. (2009). Enhancing well-being and alleviating depressive symptoms with positive psychology interventions: A practice-friendly meta- analysis. *Journal of Clinical Psychology*, 65(5), 467-487.

Sinclair, V. G., & Wallston, K. A. (2004). The development and psychometric evaluation of the Brief Resilient Coping Scale. *Assessment*, 11(1), 94-101.

Smith, L. E., Seltzer, M. M., Tager-Flusberg, H., Greenberg, J. S., & Carter, A. S. (2008). A comparative analysis of well-being and coping among mothers of toddlers and mothers of adolescents with ASD. *Journal of Autism and Developmental Disorders*,

38(5), 876-889.

Spielberger, C. D. (1983). Manual for the State-Trait Anxiety Inventory STAI (form Y)("self-evaluation questionnaire").

Tang, S. T., Chang, W. C., Chen, J. S., Wang, H. M., Shen, W. C., Li, C. Y., & Liao, Y. C. (2013). Course and predictors of depressive symptoms among family caregivers of terminally ill cancer patients until their death. *Psycho-Oncology*, 22(6), 1312-1318.

Tedeschi, R. G., & Calhoun, L. G. (1996). The Posttraumatic Growth Inventory: Measuring the positive legacy of trauma. *Journal of Traumatic Stress*, 9(3), 455-471.

Totsika, V., Hastings, R. P., Emerson, E., Lancaster, G. A., & Berridge, D. M. (2011). A population-based investigation of behavioural and emotional problems and maternal mental health: Associations with autism spectrum disorder and intellectual disability. *Journal of Child Psychology and Psychiatry*, 52(1), 91-99.

Walsh, F. (1998). *Strengthening Family Resilience*. The Guilford Family Therapy Series.

Walsh, F. (2006). *Strengthening family resilience*. Guilford Press.

Weiss, J. A., Cappadocia, M. C., MacMullin, J. A., Viecili, M. A., & Lunsy, Y. (2012). Psychological acceptance and empowerment as mediators of the impact of problem behaviour in children with autism spectrum disorders on parent mental health. *Autism: The International Journal of Research and Practice*, 16, 261-274.

Windle, G. (2011). What is resilience? A review and concept analysis. *Reviews in Clinical Gerontology*, 21(02), 152-169.

Zeedyk, S. M., & Blacher, J. (2015). Chapter One-Maternal Depression and Child Behavior

Problems: Longitudinal Considerations. *International Review of Research in  
Developmental Disabilities*, 49, 1-43.

Zigmond, A. S., & Snaith, R. P. (1983). The hospital anxiety and depression scale. *Acta  
Psychiatrica Scandinavica*, 67(6), 361-370.

Author Note

Elizabeth Halstead, Bangor University, UK

Naomi Ekas, Texas Christian University, USA

Richard P. Hastings, University of Warwick, UK

Gemma M. Griffith, Bangor University, UK

Study 1 was supported in part by a PhD scholarship to Elizabeth Halstead funded by Bangor University. Study 2 was supported by the Research and Creative Activities Fund at Texas Christian University. Portions of these data served as part of the doctoral dissertation of the first author.

Correspondence concerning this article should be addressed to Naomi Ekas, Texas Christian University, TCU Box 298920, Fort Worth, TX 76129, 817-257-7848, [naomi.ekas@tcu.edu](mailto:naomi.ekas@tcu.edu)